

Erratum

Evidence that Armadillo Transduces Wingless by Mediating Nuclear Export or Cytosolic Activation of Pangolin

Cell (2002), Vol. 111, pp. 265–280

Figures 2A–2D of this paper document the subcellular distributions of Flu-tagged, N-terminally truncated forms of Armadillo (Flu-ΔArm) that are targeted to the membrane ([A]; M⁺N⁻ Flu-ΔArm), the nucleus ([D]; M⁻N⁺ Flu-ΔArm), both ([B]; M⁺N⁺ Flu-ΔArm) or neither ([C]; M⁻N⁻ Flu-ΔArm). However, the image shown in (C) is, inadvertently, a differently cropped and rotated version of the image shown in (B). Here, we show a repeat of this experiment. As initially reported, and now correctly documented, both dual and untargeted Flu-ΔArm accumulate in the cytosol as well as the nucleus, in contrast to membrane- and nuclear-targeted Flu-ΔArm which localize predominantly outside or inside nuclei.

We take this opportunity to correct two other minor errors, both in the legend to Figure 1. First, the C terminally truncated mutant protein Arm^{XM19} is described as lacking part of the thirteenth Arm repeat as well as the C-terminal domain, a description based on the initial classification of Arm repeats by Peifer et al. (1990). However, Arm is now considered to have only twelve Arm repeats, and the Arm^{XM19} protein is truncated after the twelfth repeat. Second, ArmΔ is an engineered form of Arm which is nine, rather than eleven, amino acids longer than Arm^{XM19}.

The corrected Figure 2 and its legend are below.

S.K. Chan and G. Struhl




Figure 2. Distribution of Membrane, Nuclear, Dual, or Untargeted ΔArm Proteins. Subcellular distribution of Flu-tagged ΔArm (Green) expressed under C765-Gal4/UAS control in imaginal wing disc cells counterstained for Lamin (Red), which marks the nuclear membrane. Membrane (A; M⁺N⁻) and nuclear (D; M⁻N⁺) targeted forms accumulate predominantly outside or inside nuclei; dual targeted (B; M⁺N⁺) and untargeted (C; M⁻N⁻) forms distribute more uniformly.